

Optimal State Transfer and Entanglement Generation in Power-law Interacting Systems


(arXiv:2010.02930)

Minh Tran


Joint work with Abhinav Deshpande, Andrew Guo, Andrew Lucas, and Alexey Gorshkov

QuICS/JQI, University of Maryland


QIP, 02/02/2021



Most physical interactions decay with distance, e.g. as a power law




e.g. Magnetic atoms in optical lattice




Lepoutre et al, Nat Commun 10, 1714 (2019)

e.g. Rydberg-Rydberg interaction




Omran et al, Science 365, 570 (2019)



Coulomb


dipole-dipole

van der Waals




Zhang et al, Nature 551, 601–604 (2017)

Decaying interactions



Locality:

It takes time for information to propagate in the system



Larger distance r




Takes longer time to propagate

In general:


$$t \geq f(r)$$



Some function




Longer-range interactions



Implications of locality?

Suppose we want to apply quantum gates
between distant particles

- With nearest-neighbor interactions:
 $\Rightarrow$ Take time $t \sim r$ (time unit)
- With power-law ($1/r^\alpha$) interactions:
Can we do faster than $t \sim r$?




Transfer quantum states
Generate entanglement
Scramble information
...

Outline

- ❖ Theoretical limits: Lieb-Robinson bounds
- ❖ Optimal state transfer and entanglement generation
 - Key implications of the protocol

Setup:



- Power-law interacting Hamiltonian $H(t) = \sum_{i,j} h_{ij}(t)$ s.t. $\|h_{ij}(t)\| \leq 1/r_{ij}^\alpha \quad \forall t$
- Two operators A, B initially supported on sites separated by distance r

Lieb-Robinson bounds:


$$\| [A(t), B(0)] \| \leq ?$$

$\uparrow$ $e^{iHt} A e^{-iHt}$

$t = 0 \Rightarrow [A, B] = 0$
 $t > 0 \Rightarrow [A(t), B] \neq 0$

Because $A(t)$ spreads to B

Setup:




- Power-law interacting Hamiltonian $H(t) = \sum_{i,j} h_{ij}(t)$ s.t. $\|h_{ij}(t)\| \leq 1/r_{ij}^\alpha \quad \forall t$
- Two operators A, B initially supported on sites separated by distance r

E.g.: Nearest-neighbor ($\alpha \rightarrow \infty$)

Lieb & Robinson, CMP '72

$$\| [A(t), B(0)] \| \lesssim e^{vt-r}$$


$\Rightarrow$ Linear light cone: $t \gtrsim r$



Theoretical limits: Lieb-Robinson bounds

(d -dimensional lattice)

Related works: Hazzard et al, 2013; Hauke & Tagliacozzo, 2013; Schachenmayer et al, 2013; Knap et al, 2013; Juenemann et al, 2013; Eisert et al, 2013; Hazzard et al, 2014; Gong et al, 2014; Foss-Feig et al, 2015; Storch et al, 2015; Rajabpour et al, 2014, 2015, ...




Theoretical limits: Lieb-Robinson bounds

(d -dimensional lattice)


Related works: Hazzard et al, 2013; Hauke & Tagliacozzo, 2013; Schachenmayer et al, 2013; Knap et al, 2013; Juenemann et al, 2013; Eisert et al, 2013; Hazzard et al, 2014; Gong et al, 2014; Foss-Feig et al, 2015; Storch et al, 2015; Rajabpour et al, 2014, 2015, ...

Are the bounds tight?




Quantum state transfer


Initial state:




Target state:



Fast state-transfer protocols (d dimensions)




Fast state-transfer protocols (d dimensions)




Optimal quantum state transfer

Initial state:

$$a|0\rangle + b|1\rangle$$






Target state:



Strategy:


Encode information into r -qubit
Greenberger–Horne–Zeiling
(GHZ) state



Decode the information
into the target qubit

Encoding information into a GHZ state

Suppose we can “merge” several GHZ states into one big GHZ state




Prepare “one-qubit GHZ states”:

$$\bullet \longrightarrow |\psi\rangle = a|0\rangle + b|1\rangle$$

$$\bullet \longrightarrow |+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$$


Our strategy: Repeatedly merge nearby GHZ states into larger and larger GHZ states with the encoded information




Optimal quantum state transfer

Initial state:

$$a|0\rangle + b|1\rangle$$






Target state:



Strategy:


Encode information into r -qubit
Greenberger–Horne–Zeiling
(GHZ) state



Decode the information
into the target qubit

Subroutine: merging GHZ states


Apply $H = \frac{1}{r_{\max}^\alpha} \sum_{i,j} |1\rangle\langle 1|_i \otimes |1\rangle\langle 1|_j$



Total time $\sim \frac{r_{\max}^\alpha}{V^2}$

After time $t = \pi \frac{r_{\max}^\alpha}{V^2}$

$V = 3 =$ Volume of each small GHZ states



$$+ \begin{matrix} a & |0\rangle & |0\rangle & |0\rangle \\ b & |1\rangle & |1\rangle & |1\rangle \end{matrix} \begin{pmatrix} + & |0\rangle & |0\rangle & |0\rangle \\ & |1\rangle & |1\rangle & |1\rangle \end{pmatrix}$$

Unentangle



$$+ \begin{matrix} a & |0\rangle & |0\rangle & |0\rangle & |0\rangle & |0\rangle \\ b & |1\rangle & |1\rangle & |1\rangle & |0\rangle & |0\rangle \end{matrix} \begin{pmatrix} + & |0\rangle \\ & |1\rangle \end{pmatrix}$$

Single-qubit rotation



$$+ \begin{matrix} a & |0\rangle & |0\rangle & |0\rangle & |0\rangle & |0\rangle \\ b & |1\rangle & |1\rangle & |1\rangle & |0\rangle & |0\rangle \end{matrix}$$

Re-entangle



$$+ \begin{matrix} a & |0\rangle & |0\rangle & |0\rangle & |0\rangle & |0\rangle \\ b & |1\rangle & |1\rangle & |1\rangle & |1\rangle & |1\rangle \end{matrix}$$


Encoding information into a GHZ state

Prepare “one-qubit GHZ states”:

$$\bullet \longrightarrow |\psi\rangle = a|0\rangle + b|1\rangle$$

$$\bullet \longrightarrow |+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$$

Time for implementing
each iteration $\sim \frac{r_{\max}^\alpha}{V^2}$



Larger $V \Rightarrow$ Allow larger $r_{\max}$

$\Rightarrow$ Bigger and bigger GHZ states

Total time (asymptotically):


$$t \sim \begin{cases} r^{\alpha-2d} & (2d < \alpha < 2d+1) \\ \log^\kappa r & (\alpha < 2d) \end{cases}$$

Strategy: Repeatedly merge nearby GHZ states into larger and larger GHZ states with the encoded information

More details in

Tran et al, arXiv:2010.02930 (2020)

② Saturates* the Lieb-Robinson bound



③ **State-dependent** state transfer is provably exponentially faster than **state-independent** state transfer

- ① Subpolynomial state-transfer time for
- Trapped ions ($\alpha \in [0,3]$) with $d = 1, 2$
 - Dipole-dipole interactions ($\alpha = 3$) with $d = 2, 3$
 - Van-der-Waals interactions ($\alpha = 6$) with $d = 3$

- We need $\Omega(n)$ gates to make an n -qubit GHZ state
- Simulating $e^{-i H t}$ for $t = t_*$ must also take $\Omega(n)$ gates
- Meanwhile, Suzuki-Trotter takes $O(n^{2+o(1)})$
- Either $\exists$ better simulation algorithms or making GHZ states is strictly easier than simulation

④ A lower bound on the gate count of quantum simulation algorithms

Acknowledgments



Abhinav Deshpande
(UMD)



Andrew Guo
(UMD)



Andrew Lucas
(UC Boulder)



Alexey Gorshkov
(UMD)



JOINT CENTER FOR
QUANTUM INFORMATION
AND COMPUTER SCIENCE



JOINT QUANTUM
INSTITUTE




University of Colorado **Boulder**



UNIVERSITY OF
MARYLAND

② Saturates* the Lieb-Robinson bound



③ **State-dependent** state transfer is provably exponentially faster than **state-independent** state transfer

- ① Subpolynomial state-transfer time for
- Trapped ions ($\alpha \in [0,3]$) with $d = 1, 2$
 - Dipole-dipole interactions ($\alpha = 3$) with $d = 2, 3$
 - Van-der-Waals interactions ($\alpha = 6$) with $d = 3$

④ A lower bound on the gate count of quantum simulation algorithms

Thank you!